



**94-0091**

The use: In it's capacity of inhibitory of the agiotensine hormone. The essence: the product with a general formula:



where R<sub>1</sub> represents COOH or the group



R<sub>2</sub> represents n-propyl or n-butyl,

R<sub>3</sub> CZ, CF<sub>3</sub>, C<sub>2</sub>F<sub>5</sub>, C<sub>6</sub>H<sub>5</sub> or COOH;


R<sub>4</sub> represents COOH; or CHO or CH<sub>2</sub>OH with the condition that then R<sub>4</sub> represents CH<sub>2</sub>OH, R<sub>3</sub> represents C<sub>2</sub>F<sub>5</sub> and R<sub>2</sub> represents n-propyl;

(b) when R<sub>3</sub> represents COOH, R<sub>4</sub> is also COOH;

(c) when R<sub>2</sub> represents n-propyl, R<sub>3</sub> represents C<sub>2</sub>F<sub>5</sub> and R<sub>4</sub> represents COOH, R<sub>1</sub> constitutes the group




Reagent 1: Product with formula




where R<sub>2</sub> has the indicated values: R<sub>5</sub> represents CZ, CF<sub>3</sub>, C<sub>2</sub>F<sub>5</sub>, C<sub>6</sub>H<sub>5</sub> or COO (C<sub>1-4</sub> - alkyl); R<sub>6</sub> represents CH<sub>2</sub>OH, CO<sub>2</sub> (C<sub>1-4</sub>-alkyl) or CHO).

Reagent 2: Product with formula



where X represents halogen, n represents toluene sylphonix or methylsulphonilox; R<sub>7</sub> represents COO (C<sub>1-4</sub> - alkyl), CN or the group



The temperature of the dissolvent. A base is the catalyst.